

10/692,827

Listing of the Claims:

In The Claims:


Please amend the claims as follows.

1. (currently amended) A catalyst system comprising the reaction product of:
 - (a) an activator; and
 - (b) a catalyst precursor comprising:
 - (i) a Group-8, -9, or -10 transition metal, M;
 - (ii) an ancillary ligand comprising:
 - a terminal amine comprising two independently selected hydrocarbyl radicals, R¹ and R²
 - a terminal phosphine comprising two independently selected hydrocarbyl radicals, R³ and R⁴ and
 - a hydrocarbyl bridge, Y, comprising a backbone wherein the hydrocarbyl bridge connects between the terminal amine and the terminal phosphine and wherein the backbone comprises a chain that is four or more carbon atoms long; and
 - (iii) an abstractable ligand, X₁

wherein the activator is an alumoxane, an aluminum alkyl, an alkyl aluminum halide, an alkylaluminum alkoxide, a discrete ionic activator, or a Lewis acid.
2. (original) The catalyst system of Claim 1 wherein the catalyst precursor has the following formula:

BEST AVAILABLE COPY

10/692,827



wherein

- (i) M is a Group-8, -9, or -10 transition metal;
 - (ii) N is nitrogen;
 - (iii) P is phosphorus;
 - (iv) R¹, R², R³, and R⁴ are independently hydrocarbyl radicals;
 - (v) Y is a hydrocarbyl bridge comprising a backbone wherein the backbone comprises a chain that is four or more carbon atoms long;
 - (vi) X are independently abstractable ligands.
3. (original) A catalyst system comprising the reaction product of
 - (a) the catalyst system of Claim 2 and
 - (b) ethylene, propylene, 1-butene, or a mixture of any two or all three of ethylene, propylene, and 1-butene.
 4. (original) The catalyst system of Claim 2 further comprising at least one additional olefin polymerization catalyst.
 5. (currently amended) The catalyst system of Claim 2 wherein R¹, R², R³, and R⁴ are independently selected from C₁-C₄₀ hydrocarbyls.

10/692,827

6. (currently amended) The catalyst system of Claim 5 wherein R^1 , R^2 , R^3 , and R^4 are independently selected from C_1 - C_{30} hydrocarbyls.
7. (currently amended) The catalyst system of Claim 6 wherein R^1 , R^2 , R^3 , and R^4 are independently selected from methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, undecynyl, dodecynyl, phenyl, benzyl, phenethyl, tolyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, or cyclododecyl radicals.
8. (currently amended) The catalyst system of Claim 7 wherein R^1 , R^2 , R^3 , and R^4 are independently selected from methyl, ethyl, propyl, butyl, cyclohexyl, phenyl, tolyl, benzyl, and or phenethyl.
9. (currently amended) The catalyst system of Claim 2 wherein X are independently hydride radicals; hydrocarbyl radicals; ~~hydrocarbyl-substituted~~, hydrocarbyl-substituted; or organometalloid radicals.
10. (currently amended) The catalyst system of Claim 9 wherein two X's are connected to each other to form a 3-to-50-atom metallacycle ring.
11. (original) The catalyst system of Claim 2 wherein X are independently halogen, alkoxide, aryloxy, amide, or phosphide radicals.
12. (currently amended) The catalyst system of Claim 11 wherein X are independently chloride, bromide, iodide, methoxide, ethoxide,

I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827

dimethylamide, diethylethoxide, or phenoxide.







13. (currently amended) The catalyst system of Claim 9 wherein X are independently selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, hydride, phenyl, benzyl, phenethyl, tolyl, methoxy, ethoxy, propoxy, butoxy, dimethylamino, diethylamino, ~~methylethylamine,~~ and methylethylamino.
14. (original) The catalyst system of Claim 2 wherein X are independently acetylacetonate, 1,1, 1,5,5,5-hexa-fluoroacetylacetonate, 1,1,1-trifluoroacetylacetonate, or 1,1, 1-trifluoro-5,5-di-methylacetylacetonate radicals.
15. (currently amended) The catalyst system of Claim 2 wherein M is selected from the group consisting of nickel, iron, cobalt, palladium, platinum, ruthenium, osmium, rhodium, and iridium.
16. (currently amended) The catalyst system Claim 15 wherein M is selected from the group consisting of iron, nickel, cobalt, and palladium.
17. (currently amended) The catalyst system of Claim 15 wherein Y is selected from the group consisting of butylene, pentylene, hexylene, heptylene, octylene, nonylene, decylene, undecylene, dodecylene, tridecylene, tetradecylene, pentadecylene, hexadecylene, heptadecylene, octadecylene, nonadecylene, eicosylene, heneicosylene, docosylene, tricosylene, tetracosylene, pentacosylene, hexacosylene, heptacosylene, octacosylene, nonacosylene, triacontylene, cyclohexylene, cyclooctylene, cyclododecylene, cyclododecylene, biphenyl, butenylene, pentenylene, hexenylene, heptenylene, octenylene, nonenylene, decenylene,

I:\Bpe\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc





10/692,827

undecenylene, dodecenylene, hexynylene, heptynylene, octynylene, nonynylene, decynylene, undecynylene, dodecynylene, butadienylene, pentadienylene, hexadienylene, heptadienylene, octadienylene, nonadienylene, decadienylene, undecadienylene, dodecadienylene, hexatrienylene, octatrienylene, decatrienylene, and dodecatrienylene radicals.









18. (currently amended) The catalyst system of claim 17 wherein Y is selected from biphenyl.
19. (currently amended) The catalyst system of claim 2 wherein Y has one of the following formulas:



10/692,827



10/692,827



where


- (a) R' are independently[,] hydrogen or C_1 - C_{50} hydrocarbyl radicals;
- (b) A is a non-hydrocarbon atom functional group;
- (c) E is a Group-14 element;

I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US2002B130A-2-US-2006Feb27-1.111 Amend.doc


10/692,827

- (d) x is an integer from 1 to 4; and
 - (e) y is an integer from 0 to 4.
20. (currently amended) The catalyst system of Claim 19 wherein A is selected from the group consisting of C=O, C=S, O, S, SO₂, NR*, PR*, BR*, SiR*₂, and GeR*₂, where R* is independently a hydrocarbyl or halocarbyl radical.
21. (withdrawn) A polymerization method comprising the step of providing at least one catalyst system of Claim 2.
22. (withdrawn) The polymerization method of Claim 21 wherein the catalyst's activity exceeds 8000 moles of ethylene per mole transition metal per hour.
23. (withdrawn) The polymerization method of Claim 22 further comprising recovering a product comprising greater than 50 mol% of linear C4-C14 a-olefins based on the total weight of polymerized product.
24. (withdrawn) The polymerization method of 23 wherein the product comprises greater than 80 mol% of linear C4-C14 a-olefins.
25. (withdrawn) The polymerization method of 24 wherein the product comprises greater than 50 mol% of linear C4 and C6 a-olefins.
26. (withdrawn) The polymerization method of 25 wherein the product comprises greater than 80 mol% of linear C4 and C6 a-olefins.
27. (withdrawn) A catalyst precursor with one of the following formula:

10/692,827



28. (currently amended) A catalyst system comprising the reaction product of:
- an activator; and
 - a catalyst precursor with the following formula:



wherein

- M is iron, nickel, cobalt, and palladium;
- N is nitrogen;
- P is phosphorus;
- R^1 , R^2 , R^3 , and R^4 are independently selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl,

I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc


10/692,827

heptacosyl, octacosyl, nonacosyl, triacontyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, undecynyl, dodecynyl, phenyl, benzyl, phenethyl, tolyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, and cyclododecyl radicals;

- (v) Y is a hydrocarbonyl bridge comprising a backbone wherein the backbone comprises a chain that is four or more carbon atoms long; and
- (vi) X are independently abstractable ligands,
wherein the activator is an alumoxane, an aluminum alkyl, an alkyl aluminum halide, an alkylaluminum alkoxide, a discrete ionic activator, or a Lewis acid.

29. (currently amended) A catalyst system comprising the reaction product of:

- (a) an activator; and
- (b) a catalyst precursor with the following formula:



wherein

- (i) M is ~~from~~ nickel, iron, cobalt, palladium, platinum, ruthenium, osmium, rhodium, and iridium or iridium;

I:\Bpc\LA\W\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827

- (ii) N is nitrogen;
- (iii) P is phosphorus;
- (iv) R^1 , R^2 , R^3 , and R^4 are independently selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, undecynyl, dodecynyl, phenyl, benzyl, phenethyl, tolyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, and cyclododecyl radicals;
- (v) Y is a hydrocarbyl bridge comprising a backbone wherein the backbone comprises a chain that is four or more carbon atoms long; and
- (vi) X are independently chloride, bromide, iodide, methoxide, ethoxide, dimethylamide, diethylethoxide, phenoxide, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, hydride, phenyl, benzyl, phenethyl, tolyl, methoxy, ethoxy, propoxy, butoxy, dimethylamino, diethylamino, methylethylamino, acetylacetate, 1,1, 1,5,5,5-hexa-fluoroacetylacetate, 1,1, 1-trifluoro-acetylacetate, or 1,1, 1-trifluoro-5,5-di-methylacetylacetate radicals; or two X's are connected to form a 3-to-40-atom metallacycle ring,
wherein the activator is an alumoxane, an aluminum alkyl, an alkyl aluminum halide, an alkylaluminum alkoxide, a discrete ionic


I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827

activator, or a Lewis acid.

30. (currently amended) A catalyst system comprising the reaction product of:

- (a) an activator; and
- (b) a catalyst precursor with the following formula:



wherein

- (i) M is ~~selected from~~ nickel, iron, cobalt, palladium, platinum, ruthenium, osmium, rhodium, ~~and iridium~~ or iridium;
- (ii) N is nitrogen;
- (iii) P is phosphorus;
- (iv) R¹, R², R³, and R⁴ are independently selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, undecynyl, dodecynyl, phenyl, benzyl, phenethyl, tolyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl,

I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827


- cyclodecyl, or cyclododecyl radicals;
- (v) Y is ~~selected from~~ butylene, pentylene, hexylene, heptylene, octylene, nonylene, decylene, undecylene, dodecylene, tridecylene, tetradecylene, pentadecylene, hexadecylene, heptadecylene, octadecylene, nonadecylene, eicosylene, heneicosylene, docosylene, tricosylene, tetracosylene, pentacosylene, hexacosylene, heptacosylene, octacosylene, nonacosylene, triacontylene, cyclohexylene, cyclooctylene, cyclodecylene, cyclododecylene, biphenyl, butenylene, penentylene, hexenylene, heptenylene, octenylene, nonenylene, decenylene, undecenylene, dodecenylene, hexynylene, heptynylene, octynylene, nonynylene, decynylene, undecynylene, dodecynylene, butadienylene, pentadienylene, hexadienylene, heptadienylene, octadienylene, nonadienylene, decadienylene, undecadienylene, dodecadienylene, hexatrienylene, octatrienylene, decatrienylene, and dodecatrienylene radicals or dodecatrienylene radicals; and
- (vi) X are independently chloride, bromide, iodide, methoxide, ethoxide, dimethylamide, diethylethoxide, phenoxide, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, hydride, phenyl, benzyl, phenethyl, tolyl, methoxy, ethoxy, propoxy, butoxy, dimethylamino, diethylamino, methylethylamino, acetylacetonate, 1,1, 1,5,5,5-hexafluoroacetylacetonate, 1,1, 1-trifluoro-acetylacetonate, or 1,1, 1-trifluoro-5,5-di-methylacetylacetonate radicals; or two X's are connected to each other to form a 3-to-40-atom metallacycle ring, wherein the activator is an alumoxane, an aluminum alkyl, an alkyl aluminum halide, an alkylaluminum alkoxide, a discrete ionic activator, or a Lewis acid.

I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827

31. (currently amended) A catalyst system comprising the reaction product of:







- (a) an activator; and
- (b) a catalyst precursor with the following formula:







wherein

- (i) M is a Group-8, -9, or -10 transition metal;
- (ii) N is nitrogen;
- (iii) P is phosphorus;
- (iv) R^1 , R^2 , R^3 , and R^4 ~~R^1 , R^2 , R^3 , and R^4~~ are independently hydrocarbyl radicals;
- (v) Y is represented by one of the following formulas:









10/692,827



10/692,827



10/692,827



where

- R' are independently, hydrogen or C_1 - C_{50} hydrocarbyl radicals;
- A is a non-hydrocarbon atom functional group;
- E is a Group-14 element;


I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827

- x is an integer from 1 to 4; and
 - y is an integer from 0 to 4.
- (vi) X are independently chloride, bromide, iodide, methoxide, ethoxide, dimethylamide, diethylethoxide, phenoxide, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, hydride, phenyl, benzyl, phenethyl, tolyl, methoxy, ethoxy, propoxy, butoxy, dimethylamino, diethylamino, methylethylamino, acetylacetonate, 1,1, 1,5,5,5-hexa-fluoroacetylacetonate, 1,1, 1-trifluoro-acetylacetonate, or 1,1,1-trifluoro-5,5-di-methylacetylacetonate radicals; or two X's are connected to each other to form a 3-to-40-atom metallacycle ring, wherein the activator is an alumoxane, an aluminum alkyl, an alkyl aluminum halide, an alkylaluminum alkoxide, a discrete ionic activator, or a Lewis acid.

32. (withdrawn) A polymerization method wherein the catalysts activity exceeds 8000 moles of ethylene per mole transition metal per hour comprising the step of providing at least one catalyst system comprising the reaction product of:
- (a) an activator; and
 - (b) a catalyst precursor with the following formula:

10/692,827




wherein

- (i) M is iron, nickel, cobalt, and palladium;
- (ii) N is nitrogen;
- (iii) P is phosphorus;
- (iv) R^1 , R^2 , R^3 , and R^4 are independently selected from methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, undecynyl, dodecynyl, phenyl, benzyl, phenethyl, tolyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, cyclododecyl radicals;
- (v) Y is a hydrocarbyl bridge comprising a backbone wherein the backbone comprises a chain that is four or more carbon atoms long;
- (vi) X are independently abstractable ligands.

10/692,827

33. (withdrawn) A polymerization method wherein the catalysts activity exceeds 8000 moles of ethylene per mole transition metal per hour comprising the step of providing at least one catalyst system comprising the reaction product of:

- (a) an activator; and
 (b) a catalyst precursor with the following formula:




wherein

- (i) M is iron, nickel, cobalt, and palladium, platinum, ruthenium, osmium, rhodium, and iridium;
 (ii) N is nitrogen;
 (iii) P is phosphorus;
 (iv) R¹, R², R³, and R⁴ are independently selected from methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, undecynyl, dodecynyl, phenyl, benzyl, phenethyl, tolyl, cyclobutyl, cyclopentyl,

10/692,827

- cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, cyclododecyl radicals;
- (v) Y is a hydrocarbyl bridge comprising a backbone wherein the backbone comprises a chain that is four or more carbon atoms long; and
- (vi) X are independently chloride, bromide, iodide, methoxide, ethoxide, dimethylamide, diethylethoxide, phenoxide, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, hydride, phenyl, benzyl, phenethyl, tolyl, methoxy, ethoxy, propoxy, butoxy, dimethylamino, diethylamino, methylethylamino, acetylacetonate, 1,1, 1,5,5,5-hexa-fluoroacetylacetonate, 1,1, 1-trifluoro-acetylacetonate, or 1,1, 1-trifluoro-5,5-di-methylacetylacetonate radicals; or two X's are connected to form a 3-to-40-atom metallacycle ring.
34. (withdrawn) A polymerization method wherein the catalysts activity exceeds 8000 moles of ethylene per mole transition metal per hour comprising the step of providing at least one catalyst system comprising the reaction product of:
- (a) an activator; and
- (b) a catalyst precursor with the following formula:

10/692,827



wherein

- (i) M is iron, nickel, cobalt, and palladium, platinum, ruthenium, osmium, rhodium, and iridium;
- (ii) N is nitrogen;
- (iii) P is phosphorus;
- (iv) R^1 , R^2 , R^3 , and R^4 are independently selected from methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, undecynyl, dodecynyl, phenyl, benzyl, phenethyl, tolyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, cyclododecyl radicals;
- (v) Y is selected from butylene, pentylene, hexylene, heptylene, octylene, nonylene, decylene, undecylene, dodecylene, tridecylene, tetradecylene, pentadecylene, hexadecylene, heptadecylene, octadecylene, nonadecylene, eicosylene, heneicosylene,


10/692,827

docosylene, tricosylene, tetracosylene, pentacosylene, hexacosylene, heptacosylene, octacosylene, nonacosylene, triacontylene, cyclohexylene, cyclooctylene, cyclodecylene, cyclododecylene, biphenyl, butenylene, penentylene, hexenylene, heptenylene, octenylene, nonenylene, decenylene, undecenylene, dodecenylene, hexynylene, heptynylene, octynylene, nonynylene, decynylene, undecynylene, dodecynylene, butadienylene, pentadienylene, hexadienylene, heptadienylene, octadienylene, nonadienylene, decadienylene, undecadienylene, dodecadienylene, hexatrienylene, octatrienylene, decatrienylene, and dodecatrienylene radicals; and

- (vi) X are independently chloride, bromide, iodide, methoxide, ethoxide, dimethylamide, diethylethoxide, phenoxide, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, hydride, phenyl, benzyl, phenethyl, tolyl, methoxy, ethoxy, propoxy, butoxy, dimethylamino, diethylamino, methylethylamino, acetylacetonate, 1,1, 1,5,5,5-hexafluoroacetylacetonate, 1,1, 1-trifluoro-acetylacetonate, or 1,1, 1-trifluoro-5,5-di-methylacetylacetonate radicals; or two X's are connected to form a 3-to-40-atom metallacycle ring.

35. (withdrawn) A polymerization method wherein the catalysts activity exceeds 8000 moles of ethylene per mole transition metal per hour comprising the step of providing at least one catalyst system comprising the reaction product of:
- (a) an activator; and
 - (b) a catalyst precursor with the following formula:







10/692,827







wherein

- (i) M is a Group-8, -9, or -10 transition metal;
- (ii) N is nitrogen;
- (iii) P is phosphorus;
- (iv) R^1 , R^2 , R^3 , and R^4 are independently hydrocarbyl radicals;
- (v) Y is represented by one of the following formulas:









10/692,827



10/692,827



10/692,827



where

- R' are independently, hydrogen or C_1 - C_{50} hydrocarbyl radicals;
- A is a non-hydrocarbon atom functional group;
- E is a Group-14 element;

I:\Bpc\LA W\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827

- x is an integer from 1 to 4; and
 - y is an integer from 0 to 4.
- (vi) X are independently chloride, bromide, iodide, methoxide, ethoxide, dimethylamide, diethylethoxide, phenoxide, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl, heneicosyl, docosyl, tricosyl, tetracosyl, pentacosyl, hexacosyl, heptacosyl, octacosyl, nonacosyl, triacontyl, hydride, phenyl, benzyl, phenethyl, tolyl, methoxy, ethoxy, propoxy, butoxy, dimethylamino, diethylamino, methylethylamino, acetylacetonate, 1,1, 1,5,5,5-hexa-fluoroacetylacetonate, 1,1, 1-trifluoro-acetylacetonate, or 1,1,1-trifluoro-5,5-di-methylacetylacetonate radicals; or two X's are connected to form a 3-to-40-atom metallacycle ring.
36. (New) The catalyst system of Claim 1, wherein the activator comprises a cyclic oligomeric aluminum compound represented by the formula $(R''-Al-O)_n$, or a linear oligomeric aluminum compound represented by the formula $R''(R''-Al-O)_nAlR''_2$, wherein R'' is independently a C_1 - C_{20} alkyl radical, and wherein n is an integer from 1-50.
37. (New) The catalyst system of Claim 1, wherein the activator is methylalumoxane.
38. (New) The catalyst system of Claim 1, wherein the activator is triethylaluminum, diethylaluminum, chloride, triisobutylaluminum, tri-n-octylaluminum, or a combination thereof.
39. (New) The catalyst system of Claim 1, wherein the activator is

I:\Bpc\LA\Prosecution\EMCC Prosecution\2002\2002B130A\US\2002B130A-2-US-2006Feb27-1.111 Amend.doc

10/692,827

[Me₂PhNH][B(C₆F₅)₄], [Bu₃NH][BF₄], [NH₄][PF₆], [NH₄][SbF₆], [NH₄][AsF₆], [NH₄][B(C₆H₅)₄], B(C₆F₅)₃, B(C₆H₅)₃, or a combination thereof.

40. (New) The catalyst system of Claim 1, wherein the catalyst is deposited on a solid support, the solid support comprising polymeric materials or refractory oxide materials.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.